

Sample Size Determination in Longitudinal Studies

Don Hedeker

Department of Public Health Sciences

University of Chicago

hedeker@uchicago.edu

<https://hedeker-sites.uchicago.edu/>

Hedeker, Gibbons, & Waternaux (1999). Sample size estimation for longitudinal designs with attrition. *Journal of Educational and Behavioral Statistics*, 24:70-93

<https://hedeker-sites.uchicago.edu/page/methodological-publications-1987-1999>

1

Comparison of two groups at a single timepoint

Number of subjects (N) in each of two groups (Fleiss, 1986):

$$N = \frac{2(z_\alpha + z_\beta)^2 \sigma^2}{(\mu_1 - \mu_2)^2} = \frac{2(z_\alpha + z_\beta)^2}{[(\mu_1 - \mu_2)/\sigma]^2}$$

- z_α is the value of the standardized score cutting off $\alpha/2$ proportion of each tail of a standard normal distribution (for a two-tailed hypothesis test)
- z_β is the value of the standardized score cutting off the upper β proportion
- σ^2 is the assumed common variance in the two groups
- $\mu_1 - \mu_2$ is the difference in means of the two groups

2

Some common choices:

- $z_\alpha = 1.645, 1.96, 2.576$ for 2-tailed .10, .05, and .01 test
- $z_\beta = .842, 1.036, 1.282$ for power = .8, .85. and .90
- effect size $= (\mu_1 - \mu_2)/\sigma = .2, .5, .8$ for “small,” “medium,” and “large” effects (Cohen, 1988)

Example

- $z_\alpha = 1.96$ 2-tailed .05 hypothesis test
- $z_\beta = .842$ power = .8
- effect size $(\mu_1 - \mu_2)/\sigma = .5$

$$N = \frac{2(1.96 + .842)^2}{(.5)^2} = 15.7/.25 = 62.8$$

\Rightarrow need 63 subjects in each group

Rule of thumb: $N \approx (4/\delta)^2$, where δ = effect size
(for power = .8 for a 2-tailed .05 test)

effect size (δ)	N per group	$(4/\delta)^2$
.1	1571	1600
.2	394	400
.3	176	178
.4	100	100
.5	64	64
.6	45	44
.7	34	33
.8	26	25
.9	21	20
1.0	17	16

Amaze your friends with your sample size determination abilities!

Comparison of two groups across time

consistent difference across time

Number of subjects N in each of two groups (Diggle *et al.*, 2002)

$$N = \frac{2(z_\alpha + z_\beta)^2 (1 + (n - 1)\rho)}{n[(\mu_1 - \mu_2)/\sigma]^2}$$

- σ^2 is the assumed common variance in the two groups
- $\mu_1 - \mu_2$ is the difference in means of the two groups
- n is the number of timepoints
- ρ is the assumed correlation of the repeated measures

Example

- $z_\alpha = 1.96$ 2-tailed .05 hypothesis test
- $z_\beta = .842$ power = .8
- effect size $(\mu_1 - \mu_2)/\sigma = .5$
- $n = 2$ timepoints
- $\rho = .6$ correlation of repeated measures

$$N = \frac{2(1.96 + .842)^2(1 + (2 - 1) \times .6)}{2 \times (.5)^2} = \frac{(15.7)(1.6)}{(2)(.25)} = 50.3$$

\Rightarrow need approximately 50 subjects in each group

if $\rho = 0$ then $N = 31.4$ (cross-sectional)

if $\rho = 1$ then $N = 62.8$ (one-timepoint)

7

Stata code: longt.do

```
/* longitudinal t-test: longt.do
determines number of subjects per group
power = .8 for a 2-tailed .05 test

in the 'args' statement below:
n = number of repeated observations per subject
icc = intraclass correlation
effsize = effect size
*/
args n icc effsize
drop _all
set obs 1
gen za = invnormal(.975)
gen zb = invnormal(.80)
gen num = (2*(za + zb)^2)*(1 + ('n'-1)*'icc')
gen den = 'n'*('effsize'^2)
gen npergrp = num/den
noisily display "number of subjects per group is " %8.2f npergrp
exit
```

8

```

. cd "U:\Stata_long"
U:\Stata_long

. run longt 5 .4 .5
number of subjects per group is      32.65

. run longt 5 .5 .5
number of subjects per group is      37.67

. run longt 5 .6 .5
number of subjects per group is      42.70

```

SAS code

```

* determines number per group;
* 5 timepoints (ICC=.4);
* effect size of .5;
* power = .8 for a 2-tailed .05 test;
DATA one;
n = 5;
za = PROBIT(.975);
zb = PROBIT(.8);
rho = .4;
effsize = .5;
num = (2*(za + zb)**2)*(1 + (n-1)*rho);
den = n*(effsize**2);
npergrp = num/den;
PROC PRINT;VAR npergrp;
RUN;

```

Comparing two groups across timepoints - balanced case

As in Overall and Doyle (1994), sample size of contrast Ψ_c of group population means across n timepoints:

$$N = \frac{2(z_\alpha + z_\beta)^2 \sigma_c^2}{\Psi_c^2}$$

with

$$\Psi_c = \sum_{i=1}^n c_i (\mu_{1i} - \mu_{2i})$$

$$\sigma_c^2 = \sum_{i=1}^n c_i^2 \sigma_i^2 + 2 \sum_{i < j}^n c_i c_j \sigma_{ij}$$

- σ_i^2 = common variance in the two groups at timepoint i
- σ_{ij} = common covariance in the two groups between timepoints i and j
- c_i = contrast applied at timepoint i

If the sample size is known and the degree of power is to be determined, the formula can be re-expressed as:

$$z_\beta = \sqrt{\frac{N \Psi_c^2}{2 \sigma_c^2}} - z_\alpha = \sqrt{\frac{\Psi_c^2}{V(\hat{\Psi}_c)}} - z_\alpha$$

where the variance of the sample contrast $\hat{\Psi}_c$ equals

$$V(\hat{\Psi}_c) = \frac{2}{N} \sigma_c^2$$

Example

- $z_\alpha = 1.96$ 2-tailed .05 hypothesis test
- $z_\beta = .842$ power = .8
- $n = 2$ timepoints
- variance-covariance of repeated measures

$$V(y) = \begin{bmatrix} 1 & .6 \\ .6 & 1 \end{bmatrix}$$

13

I. Average group difference over time

- mean difference $\mu_1 - \mu_2 = .5$ at both t_1 and t_2
- time-related contrasts: $c_1 = c_2 = 1/2$ (*i.e.*, average over time)

$$\Psi_c = \frac{1}{2}(.5) + \frac{1}{2}(.5) = .5$$

$$\sigma_c^2 = \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 + 2\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)(.6) = .8$$

contrast effect size $\delta = \Psi_c / \sigma_c = .5 / \sqrt{.8} = .56$

$$N = \frac{2(1.96 + .842)^2}{(.56)^2} = 50$$

14

Notice

- if $\rho = 1$, then $\sigma_c^2 = 1$, $\delta = .5$, $N = 63$ (one-timepoint)
- if $\rho = 0$, then $\sigma_c^2 = 1/2$, $\delta = 1$, $N = 16$ (cross-sectional)

where ρ is the assumed correlation of the repeated measures

II. Group difference across time

- mean difference $\mu_1 - \mu_2 = 0$ at $t1$ and $.5$ at $t2$
- time-related contrasts: $c_1 = -1$ and $c_2 = 1$

$$\Psi_c = -1(0) + 1(.5) = .5$$

$$\sigma_c^2 = (-1)^2(1)^2 + (1)^2(1)^2 + 2(-1)(1)(.6) = .8$$

contrast effect size $\delta = \Psi_c / \sigma_c = .5 / \sqrt{.8} = .56$

$$N = \frac{2(1.96 + .842)^2}{(.56)^2} = 50$$

Notice

- if N was calculated based on t_2 only, then $N = 63$

$$H_0 : \mu_{12} = \mu_{22} \neq H_0 : (\mu_{12} - \mu_{11}) = (\mu_{22} - \mu_{21})$$

- if $\rho = 1$, then $\sigma_c^2 = 0$
- if $\rho = .9$, then $\sigma_c^2 = .2$, $\delta = 1.12$, $N = 14$
- if $\rho = 0$, then $\sigma_c^2 = 1/2$, $\delta = .25$, $N = 63$ cross-sectional

For average group effect over time

- as $\rho \uparrow$, then $N \uparrow$

since it's a between-subjects comparison of averages

\Rightarrow less subjects needed if the averages are based on more independent data

For group difference across time

- as $\rho \uparrow$, then $N \downarrow$

since it's a between-subjects comparison of a within-subjects comparison

\Rightarrow less subjects needed if the subject differences (i.e., pre to post) are based on more reliable data

More than 2 timepoints

- mean differences across time
- var-covar and/or correlation of repeated measures
- time-related contrast

3 timepoints

$t1$	$t2$	$t3$	
$1/3$	$1/3$	$1/3$	average across time
-1	0	1	linear trend
1	-2	1	quadratic trend

trend coefficients from tables of orthogonal polynomials

4 timepoints

$t1$	$t2$	$t3$	$t4$	
$1/4$	$1/4$	$1/4$	$1/4$	average across time
-3	-1	1	3	linear trend
1	-1	-1	1	quadratic trend
-1	3	-3	1	cubic trend

often investigators expect

- overall group difference, or
- group by (approximately) linear time interaction

Stata code: longt_contrast.do

```
/* determines number of subjects per group
   3 timepoints
   linear increasing effect sizes of 0 .25 .5
   group by linear contrast across time
   AR1 structure with rho=.5
   power = .8 for a 2-tailed .05 test
*/
mata
za = invnormal(.975)
zb = invnormal(.80)
meandiff = (0, .25, .5)
contrast = (-1, 0, 1)
corrmat = ( 1 , .5 , .25 \
            .5 , 1 , .5 \
            .25 , .5 , 1 )
contdiff = contrast * meandiff'
contvar = contrast * corrmat * contrast'
NperGrp = ((2*(za+zb)^2) * contvar) / (contdiff^2)
NperGrp
end
```

21

SAS IML code

```
* determines number per group;
* 3 timepoints;
* linear increasing effect sizes of 0 .25 .5;
* group by linear contrast across time;
* AR1 structure with rho=.5;
* power = .8 for a 2-tailed .05 test;
PROC IML;
za = PROBIT(.975);
zb = PROBIT(.8);
meandiff = {0, .25, .5};
contrast = {-1, 0, 1};
corrmat = { 1 .5 .25 ,
            .5 1 .5 ,
            .25 .5 1 };
contdiff = T(contrast) * meandiff;
contvar = T(contrast)*corrmat*contrast;
NperGrp = ((2*(za+zb)**2) * contvar)/(contdiff**2);
PRINT NperGrp;
```

22

What about Attrition?

- could use N from calculations as N for last timepoint
 - e.g., $N = 50$, retention at last timepoint = .9
 \Rightarrow start the study with $50/.9 = 56$ subjects
- can build the retention rate information into the sample size formula
 - Hedeker, Gibbons, & Waternaux (1999), *JEBS*, 24:70-93
program RMASS2 available at

<https://hedeker-sites.uchicago.edu/page/methodological-publications-1987-1999>
(named RMASS2.txt - need to download it and then rename it as RMASS2.exe)

23

Comparing two groups across timepoints

unbalanced case

Denote sample size in first group as N_{1i} and second group as N_{2i} at timepoint i ($i = 1, \dots, n$). The variance of the sample contrast $\hat{\Psi}_c$ equals

$$\begin{aligned} V(\hat{\Psi}_c) = & \sum_{i=1}^n c_i^2 \sigma_i^2 \left(\frac{1}{N_{1i}} + \frac{1}{N_{2i}} \right) \\ & + 2 \sum_{i < j}^n c_i c_j \sigma_{ij} \left(\frac{1}{\sqrt{N_{1i} N_{1j}}} + \frac{1}{\sqrt{N_{2i} N_{2j}}} \right) \end{aligned}$$

Notice, that if $N_{1i} = N_{2i} = N$, then

$$V(\hat{\Psi}_c) = \frac{2}{N} \sum_{i=1}^n c_i^2 \sigma_i^2 + 2 \sum_{i < j}^n c_i c_j \sigma_{ij} \quad \text{as before}$$

24

The current formulation is fine to calculate power, given the varying group sample sizes across time, for the sample contrast:

$$z_\beta = \sqrt{\frac{\Psi_c^2}{V(\hat{\Psi}_c)}} - z_\alpha$$

However, to figure out the necessary group sample sizes given power, more work is needed since these (N_{1i} and N_{2i}) vary across time in the equation for $V(\hat{\Psi}_c)$:

$$\begin{aligned} V(\hat{\Psi}_c) = & \sum_{i=1}^n c_i^2 \sigma_i^2 \left(\frac{1}{N_{1i}} + \frac{1}{N_{2i}} \right) \\ & + 2 \sum_{i < j}^n c_i c_j \sigma_{ij} \left(\frac{1}{\sqrt{N_{1i} N_{1j}}} + \frac{1}{\sqrt{N_{2i} N_{2j}}} \right) \end{aligned}$$

25

Use sample size in first group at first timepoint (N_{11}) as a reference

- define retention rates for this group as r_{1i} for timepoints $i = 1, \dots, n$, which indicate the proportion of N_1 subjects observed at timepoint i
(note that $r_{11} = 1$ and $N_{1i} = r_{1i} N_{11}$)
- similarly, define N_{21} and r_{2i} for group two

Then,

$$\begin{aligned} V(\hat{\Psi}_c) = & \frac{1}{N_{11}} \left[\sum_{i=1}^n c_i^2 \sigma_i^2 \left(\frac{1}{r_{1i}} + \frac{1}{r_{2i}} \frac{N_{11}}{N_{21}} \right) \right. \\ & \left. + 2 \sum_{i < j}^n c_i c_j \sigma_{ij} \left(\frac{1}{\sqrt{r_{1i} r_{1j}}} + \frac{N_{11}}{N_{21}} \frac{1}{\sqrt{r_{2i} r_{2j}}} \right) \right] \end{aligned}$$

26

and, denoting the ratio of sample sizes at the first timepoint (N_{11}/N_{21}) as $N_{.1}$, then

$$V(\hat{\Psi}_c) = \frac{1}{N_{11}} \left[\sum_{i=1}^n c_i^2 \sigma_i^2 \left(\frac{1}{r_{1i}} + \frac{N_{.1}}{r_{2i}} \right) + 2 \sum_{i < j}^n c_i c_j \sigma_{ij} \left(\frac{1}{\sqrt{r_{1i} r_{1j}}} + N_{.1} \frac{1}{\sqrt{r_{2i} r_{2j}}} \right) \right]$$

27

If the retention rates are equal for the two groups across time $r_{1i} = r_{2i} = r_i$, then

$$\begin{aligned} V(\hat{\Psi}_c) &= \frac{N_{.1} + 1}{N_{11}} \left[\sum_{i=1}^n \frac{c_i^2 \sigma_i^2}{r_i} + 2 \sum_{i < j}^n \frac{c_i c_j \sigma_{ij}}{\sqrt{r_i r_j}} \right] \\ &= \frac{N_{.1} + 1}{N_{11}} \sigma_{rc}^2 \end{aligned}$$

where σ_{rc}^2 extends σ_c^2 given earlier, namely

$$\sigma_c^2 = \sum_{i=1}^n c_i^2 \sigma_i^2 + 2 \sum_{i < j}^n c_i c_j \sigma_{ij}$$

for the case where sample sizes vary across timepoints (although group retention rates are assumed equal)

28

To calculate power for any of the above variance formulations of the sample contrast,

$$z_\beta = \sqrt{\frac{\Psi_c^2}{V(\hat{\Psi}_c)}} - z_\alpha$$

In particular, for the case of common retention rates across time

$$z_\beta = \sqrt{\left(\frac{N_{11}}{N_{.1} + 1}\right) \frac{\Psi_c^2}{\sigma_{rc}^2}} - z_\alpha$$

where $N_{.1}$ is the sample size ratio between groups

Re-expressing, the number of subjects needed in the first group at the first timepoint equals:

$$N_{11} = \frac{(N_{.1} + 1)(z_\alpha + z_\beta)^2 \sigma_{rc}^2}{\Psi_c^2}$$

Based on

- sample size ratio between groups $N_{.1}$ at the first timepoint
- equal retention rates r_i across time

\Rightarrow required sample size at each timepoint for both groups can be calculated in a relatively simple way

RMASS2: Repeated Measures with Attrition: Sample Sizes for 2 Groups

Donald Hedeker and Suna Barlas

- Calculates sample size for a 2-group repeated measures design
- Allows for attrition and a variety of variance-covariance structures for the repeated measures
- Details on the methods can be found in Hedeker, Gibbons, and Waternaux (1999, *Journal of Educational and Behavioral Statistics*, 24:70-93)
- Program runs at the “Command Prompt” and the user is queried for program parameters
- For each query, the default parameter value is given in []; hitting a carriage return sets the parameter to the default
- download RMASS2.EXE and RMASS2.PDF at <https://hedeker-sites.uchicago.edu/page/methodological-publications-1987-1999>

31

Program Parameters

fout - output file name

n - number of timepoints (maximum is 20)

alpha - alpha level for statistical test (possible values = .01, .05, .10)

nside - sided test (1 or 2)

beta - level of power (from .5 to .95 in multiples of .05)

ratio - ratio of sample sizes (group 1 to group 2)

attrit - attrition across time (1=yes, 2=no)

- *if attrit=1* - attrition rates between adjacent timepoints (assumed equal for both groups)

mtype - type of expected group differences (0=means, 1=effect sizes)

- *if mtype=0* - expected difference in group means at each timepoint
- *if mtype=1* - estype - effect size type (0=constant, 1=linear trend, 2=user-defined)
 - *if estype=0* - expected effect size (equal across time)
 - *if estype=1* - expected effect size at last timepoint
 - *if estype=2* - expected effect size at each timepoint

32

vtype - variance-covariance structure of repeated measures

- if $vtype=0$ (no random effects) $\Sigma_y = \sigma_j^2 \mathbf{R}$, $j = 1, \dots, n$ timepoints
 - standard deviation at each timepoint σ_j
 - correlation structure of repeated measures (\mathbf{R} : 1=AR1; 2=toeplitz or banded matrix; 3=all correlations equal)
- if $vtype=1$ (random-effects structure) $\Sigma_y = \mathbf{X}\Sigma_v\mathbf{X}' + \sigma^2\mathbf{\Omega}$
 - nr = number of random effects (maximum is 4)
 - random-effects variance-covariance matrix Σ_v
 - random-effects design matrix \mathbf{X} ($n \times nr$ elements)
 - error variance σ^2 and autocorrelated error structure $\mathbf{\Omega}$

contrast - type of time-related contrast for statistical test
(0=average across time, 1=linear trend, 2=user-defined)

- if $contrast=2$ - contrast coefficient at each timepoint
- this selection should generally match the effect size type selected

Example

- $z_\alpha = 1.96$ 2-tailed .05 hypothesis test
- $z_\beta = .842$ power = .8
- $n = 2$ timepoints, retention rates $r_1 = 1$ and $r_2 = .8$
- sample size ratio $N_{.1} = 1$
- variance-covariance of repeated measures

$$V(y) = \begin{bmatrix} 1 & .6 \\ .6 & 1 \end{bmatrix}$$

I. Average group difference over time

- mean difference $\mu_1 - \mu_2 = .5$ at both t_1 and t_2
- time-related contrasts: $c_1 = c_2 = 1/2$

$$\Psi_c = \frac{1}{2}(.5) + \frac{1}{2}(.5) = .5$$

$$\sigma_{rc}^2 = \left(\frac{1}{2}\right)^2 + \frac{(1/2)^2}{.8} + 2\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)(.6)/\sqrt{.8} = .9$$

contrast effect size $\delta = \Psi_c/\sigma_{rc} = .5/\sqrt{.9} = .53$

$$N_{11} = \frac{2(1.96 + .842)^2}{(.53)^2} = 56.4$$

Note:if $r_2 = 1$ then $N_{11} = 50$

35

```

Command Prompt - rmass2

Enter the output file name [rmass2.out]
Enter the number of time points [ 2]
Enter the alpha level [0.05]
One or two sided test (1 OR 2) [2]
Enter the power level [0.90] .8
Enter the sample size ratio (grp 1 to grp 2) [ 1.00]
Any attrition across time? (1=yes 2=no) [2] 1
Enter the attrition rate between timepoints 1 and 2 [0.050] .2
Enter mean diffs (=0) or effect sizes (=1) [1]
Enter the type of effect size desired [0]
0 = constant across time
1 = linear trend across time
2 = user-defined
Mean difference in SD units (effect size) [0.500]
for constant: list the eff. size that is assumed equal across time
Variance-covariance structure of repeated measures
no random effects (=0) or with random effects (=1) [0]
Enter the standard deviation at timepoint 1 [ 1.000]
Enter the standard deviation at timepoint 2 [ 1.000]
Enter correlation structure of repeated measures [1]
1 = all correlations equal
2 = stationary AR1
3 = non-stationary AR1
4 = toeplitz (banded) matrix
Enter correlation term number 1 [0.500] .6
Enter the type of time-related contrast desired [0]
0 = average across time; 1 = linear trend; 2 = user-defined
Composite Effect size (without attrition) = 0.559017
N Subj for Grp1 Time 1 (without attrition) = 50.247707
Composite Effect size (adjusted for attrition) = 0.527659
N Subj for Grp1 Time 1 (adjusted for attrition)= 56.397411
Exit the program (1=Y OR 2=N) [2] ?

```

36

```

RMAS2.OUT - Notepad
File Edit Format View Help
RMAS2 - Repeated Measures with Attrition: Sample Sizes for 2 Group Designs
-----
Correlation Matrix of Y across time
      1      2
1      1.000
2      0.600 1.000

Number of Timepoints      =      2
Alpha level               = 0.050 (2-sided)
Power level               = 0.800
Grp1 to Grp2 Sample Size Ratio = 1.000

Retention rate = 1.000 0.800
Mean Diffs    = 0.500 0.500
Stand. Devs.  = 1.000 1.000
Effect Sizes   = 0.500 0.500
Contrasts     = 0.707 0.707

Composite Mean Difference      = 0.707107
Composite Variance (without attrition) = 1.600000
Composite Variance (adjusted for attrition) = 1.795820

Composite Effect size (without attrition) = 0.559017
N Subj for Grp1 at Time 1 (without attrition) = 50.247707

Composite Effect size (adjusted for attrition) = 0.527659
N Subj for Grp1 at Time 1 (adjusted for attrition) = 56.397411

Sample Sizes by Group across Time - without Attrition
Group 1      50.2      50.2
Group 2      50.2      50.2

Sample Sizes by Group across Time - with Attrition
Group 1      56.4      45.1
Group 2      56.4      45.1
-----

```

37

II. Group difference across time

- mean difference $\mu_1 - \mu_2 = 0$ at t_1 and $.5$ at t_2
- time-related contrasts: $c_1 = -1$ and $c_2 = 1$

$$\Psi_c = -1(0) + 1(.5) = .5$$

$$\sigma_{rc}^2 = (-1)^2(1)^2 + \frac{(1)^2(1)^2}{.8} + 2(-1)(1)(.6)/\sqrt{.8} = .91$$

$$\text{contrast effect size } \delta = \Psi_c / \sigma_{rc} = .5 / \sqrt{.91} = .525$$

$$N_{11} = \frac{2(1.96 + .842)^2}{(.525)^2} = 57.1$$

Note: if $r_2 = 1$ then $N_{11} = 50$

```

Command Prompt - rmass2

Enter the output file name [rmass2.out]
Enter the number of time points [ 2]
Enter the alpha level [0.05]
One or two sided test (1 OR 2) [2]
Enter the power level [0.90] .8
Enter the sample size ratio (grp 1 to grp 2) [ 1.00]
Any attrition across time? (1=yes 2=no) [2] 1
Enter the attrition rate between timepoints 1 and 2 [0.050] .2
Enter mean diffs (=0) or effect sizes (=1) [1]
Enter the type of effect size desired [0]
0 = constant across time
1 = linear trend across time
2 = user-defined 1
Mean difference in SD units (effect size) [0.500]
for linear: list the effect size at the last timepoint .5
Variance-covariance structure of repeated measures
no random effects (=0) or with random effects (=1) [0]
Enter the standard deviation at timepoint 1 [ 1.000]
Enter the standard deviation at timepoint 2 [ 1.000]
Enter correlation structure of repeated measures [1]
1 = all correlations equal
2 = stationary AR1
3 = non-stationary AR1
4 = toeplitz (banded) matrix
Enter correlation term number 1 [0.500] .6
Enter the type of time-related contrast desired [0]
0 = average across time; 1 = linear trend; 2 = user-defined 1
Composite Effect size (without attrition) = 0.559017
N Subj for Grp1 Time 1 (without attrition) = 50.247707
Composite Effect size (adjusted for attrition) = 0.524616
N Subj for Grp1 Time 1 (adjusted for attrition) = 57.053710
Exit the program (1=Y OR 2=N) [2] ?

```

39

```

RMAS2.OUT - Notepad
File Edit Format View Help
RMAS2 - Repeated Measures with Attrition: Sample Sizes for 2 Group Designs
-----
Correlation Matrix of Y across time

      1      2
1      1.000
2      0.600 1.000

Number of Timepoints      =      2
Alpha level                = 0.050 (2-sided)
Power level                = 0.800
Grp1 to Grp2 Sample Size Ratio = 1.000

Retention rate = 1.000 0.800
Mean Diffs    = 0.000 0.500
Stand. Devs.  = 1.000 1.000
Effect Sizes  = 0.000 0.500
Contrasts     = -0.707 0.707

Composite Mean Difference      = 0.353553
Composite Variance (without attrition) = 0.400000
Composite Variance (adjusted for attrition) = 0.454180

Composite Effect size (without attrition) = 0.559017
N Subj for Grp1 at Time 1 (without attrition) = 50.247707

Composite Effect size (adjusted for attrition) = 0.524616
N Subj for Grp1 at Time 1 (adjusted for attrition) = 57.053710

Sample Sizes by Group across Time - without Attrition

Group 1    50.2    50.2
Group 2    50.2    50.2

Sample Sizes by Group across Time - with Attrition

Group 1    57.1    45.6
Group 2    57.1    45.6
-----

```

40

Dichotomous outcomes

Comparison of two groups at a single timepoint

Number of subjects (N) in each of two groups (Fleiss, 1981):

$$N = \frac{[z_\alpha(2\bar{p}\bar{q})^{1/2} + z_\beta(p_1q_1 + p_2q_2)^{1/2}]^2}{(p_1 - p_2)^2}$$

- p_1 = response proportion in group 1 ($q_1 = 1 - p_1$)
- p_2 = response proportion in group 2 ($q_2 = 1 - p_2$)
- $\bar{p} = (p_1 + p_2)/2$
- $\bar{q} = 1 - \bar{p}$

41

Example

- $z_\alpha = 1.96$ 2-tailed .05 hypothesis test
- $z_\beta = .842$ power = .8
- $p_1 = .5$ and $p_2 = .7$

$$\begin{aligned} N &= \frac{[1.96(2 \times .6 \times .4)^{1/2} + .842(.5 \times .5 + .7 \times .3)^{1/2}]^2}{(.5 - .7)^2} \\ &= 93.03 \end{aligned}$$

42

Dichotomous outcomes - longitudinal case

The number of subjects (N) in each of two groups for a consistent difference in proportions $p_1 - p_2$ between two groups across n timepoints (Diggle *et al.*, (2002):

$$N = \frac{\left[z_\alpha(2\bar{p}\bar{q})^{\frac{1}{2}} + z_\beta(p_1q_1 + p_2q_2)^{\frac{1}{2}} \right]^2 (1 + (n - 1)\rho)}{n(p_1 - p_2)^2}$$

- p_1 = response proportion in group 1 ($q_1 = 1 - p_1$)
- p_2 = response proportion in group 2 ($q_2 = 1 - p_2$)
- $\bar{p} = (p_1 + p_2)/2$
- $\bar{q} = 1 - \bar{p}$
- ρ is the common correlation across the n observations

43

Example

- $z_\alpha = 1.96$ 2-tailed .05 hypothesis test
- $z_\beta = .842$ power = .8
- $n = 2$ timepoints
- correlation of repeated outcomes = .6
- $p_1 = .5$ and $p_2 = .7$

$$\begin{aligned} N &= \frac{\left[1.96(2 \times .6 \times .4)^{\frac{1}{2}} + .842(.5 \times .5 + .7 \times .3)^{\frac{1}{2}} \right]^2 (1 + (2 - 1).6)}{2(.5 - .7)^2} \\ &= 74.42 \end{aligned}$$

if $\rho = 0$ then $N = 46.51$ (cross-sectional)

if $\rho = 1$ then $N = 93.03$ (one-timepoint)

44

Stata code: longp.do

```
/* longitudinal difference in proportions test: longp.do
determines number of subjects per group
power = .8 for a 2-tailed .05 test

in the 'args' statement below:
n = number of repeated observations per subject
icc = intraclass correlation
p1 = proportion for group 1
p2 = proportion for group 2
*/
args n icc p1 p2
drop _all
set obs 1
gen za = invnormal(.975)
gen zb = invnormal(.80)
gen q1 = 1-'p1'
gen q2 = 1-'p2'
gen pbar = ('p1'+ 'p2')/2
gen qbar = (q1+q2)/2
gen num = ((za*sqrt(2*pbar*qbar) + zb*sqrt('p1'*q1 + 'p2'*q2))^2)*(1 + ('n'-1)*'icc')
gen den = 'n'*(('p1'-'p2')^2)
gen npergrp = num/den
noisily display "number of subjects per group is " %8.2f npergrp
exit
```

45

```
. cd "U:\Stata_long"
U:\Stata_long

. run longp 4 .4 .1 .2
number of subjects per group is    109.43

. run longp 4 .5 .1 .2
number of subjects per group is    124.35

. run longp 4 .6 .1 .2
number of subjects per group is    139.27
```

46

SAS code

```
* determines number per group;
* 5 timepoints (ICC=.4);
* difference in proportions of .5 and .67 (OR=2);
* power = .8 for a 2-tailed .05 test;
DATA one;
za = PROBIT(.975);
zb = PROBIT(.8);
n = 5;
p1 = .5; p2 = 2/3;
q1 = 1-p1; q2 = 1-p2;
pbar = (p1+p2)/2;
qbar = (q1+q2)/2;
rho = .4;
num = ((za*SQRT(2*pbar*qbar) + zb*SQRT(p1*q1 + p2*q2))**2)
*(1 + (n-1)*rho);
den = n*((p1-p2)**2);
npergrp = num/den;
PROC PRINT;VAR npergrp;
RUN;
```

47

Calculate Power via Simulation

- Randomly generate large number of datasets (NumDat) with assumed parameter values
 - NumDat = 5,000 and N per group = 63
 - Observations are normally distributed with $\mu_1 = 0$, $\mu_2 = .5$, $\sigma = 1$ (*e.g.*, effect size of .5)
- Analyze each dataset (5,000 *t*-tests) and count the number of times $H_0 : \mu_1 = \mu_2$ is rejected (NumRej)
- Power = NumRej / NumDat

⇒ with above specifications, power = .8018 via simulation

48

Why simulate to get power?

- For simple situations where formulas exist, no real advantage to simulation approach
- However, for not-so-simple situations, simulation comes to the rescue
 - more complicated models (can easily include covariates and interactions)
 - different kinds of outcomes (binary, ordinal, counts)
 - can deal with longitudinal and/or clustered data

49

Comparison of Pre Post models

$X_i = \text{pre}, Y_i = \text{post}, G_i = \text{group (0=control, 1=test)}$

Post t-test

$$Y_i = \beta_0 + \beta_1 G_i + \epsilon_i$$

Change score t-test

$$(Y_i - X_i) = \beta_0 + \beta_1 G_i + \epsilon_i$$

ANCOVA

$$Y_i = \beta_0 + \beta_1 G_i + \beta_2 X_i + \epsilon_i$$

$H_0 : \beta_1 = 0$ is test of interest in all cases

50

Simulation results: tests of $H_0 : \beta_1 = 0$

- 10000 datasets with 100 subjects in each of 2 groups
- mean difference of 0 at pre, .4 at post
- variance = 1 at both timepoints for both groups
- correlation = .4, .45, .5, .55, .6 between pre and post measurements

correlation	model	rejection rate
0.400	ttest	0.81
0.400	change	0.73
0.400	ancova	0.87
0.450	ttest	0.81
0.450	change	0.77
0.450	ancova	0.89
0.500	ttest	0.81
0.500	change	0.81
0.500	ancova	0.91
0.550	ttest	0.81
0.550	change	0.85
0.550	ancova	0.92
0.600	ttest	0.81
0.600	change	0.88
0.600	ancova	0.94

51

Stata simulation program: 2 sample t-test

```
/* ttestpow.do - does power estimation for 2 group t-test

based on poispow.do from
    http://www.stata.com/support/faqs/statistics/power-by-simulation/

model: mu = b0 + b1*x  where x is a dummy variable
       y ~ Normal(mu,sd)

Specifically, power is estimated for testing the hypothesis b1 = 0, against a
user-specified alternative for a user-specified sample size. Without loss of
generality, we can assume the true value of b0 is 0 and sd is 1

In the 'args' statement below:

    N is number of simulated datasets
    nobs is the number of observations in each of the two groups
    b1 is the "true" value of b1 (the alternative hypothesis)
*/

args N nobs b1
drop _all
set obs 2
generate x=0 in 1
replace x=1 in 2
expand 'nobs'
```

52

```

sort x
generate mu='b1'*x
save tempx, replace

/* Note: Here, I generated my "x" and mu-values and stored them in a dataset
-tempx- so that the same values could be used throughout the simulation */

/* set up a dataset with N observations which will eventually contain N
"p"-values obtained by testing b1=0. */
drop _all
set obs 'N'
generate pv = .

/* Loop through the simulations */

local i 0
while 'i' < 'N' {
    local i='i'+1
    preserve

    use tempx,clear          /* get the n = 2*nobs observations of x
                             and the mean mu into memory */
    gen xn = rnormal(mu,1) /* generate n obs. of a Std Normal(mu)random
                             variable in variable -xn- */
    quietly regress xn x      /* do the regression */
    matrix V=e(V)             /* get the standard-error matrix */
    matrix b=e(b)             /* get the vector of estimated coefficients */

```

53

```

    scalar tv=b[1,1]/sqrt(V[1,1])          /* the "t"-ratio */
    scalar pv = 2*(1-normal(abs(tv)))       /* the p-value */
    restore /* get back original dataset with N observations */
    quietly replace pv=scalar(pv) in 'i'     /* set pv to the p-value for
                                             the ith simulation */
    _dots 'i' 0
}

/*The dataset in memory now contains N simulated p-values. To get an
estimate of the power, say for alpha=.05, just calculate the proportion
of pv's that are less than 0.05: */

count if pv<.05
scalar power=r(N)/'N'
scalar n=2*'nobs'
noisily display "Power is " %8.4f scalar(power) " for a sample size of " /*
*/ scalar(n) " and alpha = .05"
exit

```

54

```

. cd "u:\Stata_Examples"
u:\Stata_Examples

. run ttestpow 1000 99 0.5
Power is 0.9470 for a sample size of 198 and alpha = .05

. run ttestpow 1000 99 0.4
Power is 0.7930 for a sample size of 198 and alpha = .05

. . run ttestpow 5000 99 .4
Power is 0.8010 for a sample size of 198 and alpha = .05

. run ttestpow 5000 63 .5
Power is 0.8052 for a sample size of 126 and alpha = .05

. run ttestpow 5000 45 .6
Power is 0.8064 for a sample size of 90 and alpha = .05

```

⇒ similar SAS program is `Ttest_power_N=63.sas`

Stata simulation: logistic regression with 2 groups

```

/* lregpow.do - does power estimation for 2 group logistic regression

based on poispow.do from http://www.stata.com/support/faqs/statistics/power-by-simulation/

model:  $\mu = b_0 + b_1x$  where  $x$  is a dummy variable
        $y \sim \text{Logistic}(\mu, \pi^{2/3})$ 

Specifically, power is estimated for testing the hypothesis  $b_1 = 0$ , against a
user-specified alternative for a user-specified sample size. Without loss of
generality, we can assume the true value of  $b_0$  is 0

In the 'args' statement below:

    N is number of simulated datasets
    nobs is the number of observations in each of the two groups
    oddsratio is the "true" value of  $\exp(b_1)$  (the alternative hypothesis)
*/

args N nobs oddsratio
drop _all
set obs 2
generate x=0 in 1
replace x=1 in 2
expand 'nobs'
sort x

```

```

generate mu=log('oddsratio')*x
save tempx, replace

/* Note: Here, I generated my "x" and mu-values and stored them in a dataset
-tempx- so that the same values could be used throughout the simulation */

/* set up a dataset with N observations which will eventually contain N
"p"-values obtained by testing b1=0. */
drop _all
set obs 'N'
generate pv = .

/* Loop through the simulations */

local i 0
while 'i' < 'N' {
    local i='i'+1
    preserve

    use tempx,clear          /* get the n = 2*nobs observations of x
                             and the mean mu into memory */
    gen ystar = mu - log(1/uniform() - 1) /* generate n obs. of a Std Logistic random
                                         variable in variable -xn- */
    egen y = cut(ystar), at(-10000,0,10000)
    quietly logit y x        /* do the logistic regression */
    matrix V=e(V)            /* get the standard-error matrix */
    matrix b=e(b)            /* get the vector of estimated coefficients */

```

57

```

    scalar tv=b[1,1]/sqrt(V[1,1])          /* the "t"-ratio */
    scalar pv = 2*(1-normal(abs(tv)))      /* the p-value */
    restore /* get back original dataset with N observations */
    quietly replace pv=scalar(pv) in 'i'    /* set pv to the p-value for
                                           the ith simulation */
    _dots 'i' 0
}

/*The dataset in memory now contains N simulated p-values. To get an
estimate of the power, say for alpha=.05, just calculate the proportion
of pv's that are less than 0.05: */

count if pv<.05
scalar power=r(N)/'N'
scalar n=2*'nobs'
noisily display "Power is " %8.4f scalar(power) " for a sample size of " /*
*/ scalar(n) " and alpha = .05"
exit

```

58

```

. cd "u:\Stata_Examples"
u:\Stata_Examples

. run lregpow 1000 94 2.3333
Power is 0.8070 for a sample size of 188 and alpha = .05

. run lregpow 1000 110 2
Power is 0.7120 for a sample size of 220 and alpha = .05

. run lregpow 1000 120 2
Power is 0.7480 for a sample size of 240 and alpha = .05

. run lregpow 1000 140 2
Power is 0.8210 for a sample size of 280 and alpha = .05

```

⇒ similar SAS program is `LReg_power_N=137.sas`

Stata simulation: random int model with 2 groups

```

/* longreg.do - does power estimation for 2 group random-intercept regression

based on poispow.do from http://www.stata.com/support/faqs/statistics/power-by-simulation/

model:  $\mu = b_0 + b_1x$  where x is a dummy variable
        $y \sim \text{Normal}(\mu, sd + sdu)$ 

Specifically, power is estimated for testing the hypothesis  $b_1 = 0$ , against a
user-specified alternative for a user-specified sample size. Without loss of
generality, we can assume the true value of  $b_0$  is 0 and  $sd$  is 1

In the 'args' statement below:

    N is number of simulated datasets
    nobs is the number of subjects in each of the two groups
    b1 is the "true" value of  $b_1$  (the alternative hypothesis)
    ni is the number of repeated measures per subject
    icc is the intraclass correlation
*/

args N nobs b1 ni icc
drop _all
set obs 2
generate x=0 in 1
replace x=1 in 2
expand 'nobs'
sort x

```

```

generate mu = 'b1'*x
generate varu = 'icc'
generate vare = 1-'icc'
save tempx, replace

/* Note: Here, I generated my "x" and mu-values and stored them in a dataset
-tempx- so that the same values could be used throughout the simulation */

/* set up a dataset with N observations which will eventually contain N
"p"-values obtained by testing b1=0. */
drop _all
set obs 'N'
generate pv = .

/* Loop through the simulations */

local i 0
while 'i' < 'N' {
    local i='i'+1
    preserve

    use tempx,clear          /* get the n = 2*nobs*ni observations of x
                              and the mean mu into memory */
    generate u_i = rnormal(0,sqrt(varu))
    generate mu_i = mu + u_i
    generate subject = _n
    expand 'ni'

```

61

```

generate e_ij = rnormal(0,sqrt(vare))
generate y_ij = mu_i + e_ij
sort subject
    quietly mixed y_ij x || subject:, mle /* do the multilevel regression */
    matrix V=e(V) /* get the standard-error matrix */
    matrix b=e(b) /* get the vector of estimated coefficients */
    scalar tv=b[1,1]/sqrt(V[1,1]) /* the "t"-ratio */
    scalar pv = 2*(1-normal(abs(tv))) /* the p-value */
    restore /* get back original dataset with N observations */
    quietly replace pv=scalar(pv) in 'i' /* set pv to the p-value for
                                         the ith simulation */
    _dots 'i' 0
}

/*The dataset in memory now contains N simulated p-values. To get an
estimate of the power, say for alpha=.05, just calculate the proportion
of pv's that are less than 0.05: */

count if pv<.05
scalar power=r(N)/'N'
scalar n=2*'nobs'
scalar nrep='ni'
scalar r='icc'
scalar eff='b1'
noisily display "Power is " %8.4f scalar(power) " for a sample size of " /*
*/ scalar(n) " with " scalar(nrep) " repeated obs, effect size = " scalar(eff) , " icc = " /*
*/ scalar(r) ", and alpha = .05"
exit

```

62

```
. run longreg 1000 50 .5 5 .4
```

Power is 0.9340 for a sample size of 100 with 5 repeated obs,
effect size = .5 icc = .4, and alpha = .05

- works, but slow to run in Stata
- similar programs in SAS are much faster
`RandInt_power_N=33.sas` (random intercept model)
`LR_RandInt_power_N=70.sas` (random int logistic model)
- maybe due to my lack of Stata knowledge

Additional papers, some with programs, on power for longitudinal studies

- Basagaña X. & Spiegelman D. (2010). Power and sample size calculations for longitudinal studies comparing rates of change with a time-varying exposure. *Statistics in Medicine*, 29(2):181-92.
- Comulada W.S. & Weiss R.E. (2010). Sample size and power calculations for correlations between bivariate longitudinal data. *Statistics in Medicine*, 29(27):2811-24.
- Dang, Q., Mazumdar, S. & Houck, P.R. (2008). Sample size and power calculations based on generalized linear mixed models with correlated binary outcomes. *Computer Methods and Programs in Biomedicine*, 91(2), 122-127.
- Donohue, M.C., Gamst, A.C., & Edland, S.D. (2010). longpower: Sample size calculations for longitudinal data. <http://cran.r-project.org/web/packages/longpower/index.html>
- Jung, S.H. & Ahn, C. (2003). Sample size estimation for GEE method for comparing slopes in repeated measurements data. *Statistics in Medicine*, 22(8):1305-15.
- Raudenbush, S.W., Xiao-Feng L. (2001). Effects of study duration, frequency of observation, and sample size on power in studies of group differences in polynomial change. *Psychological Methods*, 6(4):387-401.
- Rochon, J. (1998). Application of GEE procedures for sample size calculations in repeated measures experiments. *Statistics in Medicine*, 17(14):1643-1658.
- Roy, A., Bhaumik, D.K., Subhash, S. & Gibbons R.D. (2007). Sample size determination for hierarchical longitudinal designs with differential attrition rates. *Biometrics*, 63(3):699-707. <http://healthstats.org/rmass/>
- Tu, X.M., Kowalski, J., Zhang, J., Lynch, K.G., & Crits-Christoph P. (2004). Power analyses for longitudinal trials and other clustered designs. *Statistics in Medicine*, 23(18):2799-815.
- Tu, X.M., Zhang, J., Kowalski, J., Shults, J., Feng, C., Sun, W., & Tang, W. (2007). Power analyses for longitudinal study designs with missing data. *Statistics in Medicine*, 26(15):2958-81.
- Zhang, Z., & Wang, L. (2009). Power analysis for growth curve models using SAS. *Behavior Research Methods*, 41(4), 1083-1094. <http://www.psychstat.org/us>