

# OBITUARY

## Rainer (Ray) W. Guillery

### 28 August 1929–7 April 2017

S. Murray Sherman,<sup>1</sup> Carol A. Mason,<sup>2</sup> Kutay Deniz Atabay,<sup>3</sup> Jon H. Kaas,<sup>4</sup> Anthony-Samuel LaMantia,<sup>5</sup> Anna Mitchell<sup>6</sup> and Christopher Walsh<sup>7,8</sup>

<sup>1</sup>Department of Neurobiology, The University of Chicago, Chicago, IL, USA

<sup>2</sup>Departments of Pathology and Cell Biology, Neuroscience & Ophthalmology, Columbia University, New York, NY, USA

<sup>3</sup>Department of Brain and Cognitive Sciences, Whitehead Institute for Biomedical Research, Massachusetts Institute of Technology, Cambridge, MA, USA

<sup>4</sup>Department of Psychology, Vanderbilt University, Nashville, TN, USA

<sup>5</sup>Institute for Neuroscience, The George Washington School of Medicine and Health Sciences, Washington, DC, USA

<sup>6</sup>Department of Experimental Psychology, University of Oxford, Oxford, UK

<sup>7</sup>Division of Genetics and Genomics and Howard Hughes Medical Institute, Boston Children's Hospital, Boston, MA, USA

<sup>8</sup>Departments of Pediatrics and Neurology, Harvard Medical School, Boston, MA, USA



Rainer (Ray) Guillery was a major figure in the field of neuroscience. He was remarkable along many dimensions: his research output represents a highly productive and influential body of work; his work profoundly advanced our understanding of the related fields of development and thalamocortical functioning; his work was highly imaginative and insightful; and he was a cherished colleague and role model for his many former students and friends in the field.

#### Early Life

Ray was born on 28 August 1929, in Greifswald, Germany, where his father had a university appointment as a pathologist. Ray then moved with his mother to Berlin when he was three years old, soon after his parents divorced. Ray's mother was born Jewish (his father

was not), and although she became a Quaker, the Nazis considered her to be a Jew. Luckily for us, this fact suggests a certain level of drama leading to Ray's survival and his eventual significant contributions to science. These events possibly stamped on Ray a cynicism of authority, in academic life and beyond, but suffused with his humour, never hampered his scientific and professional progression and accomplishments. He left Germany with his mother in the fall of 1938, and after an odyssey that saw him spend time in the Netherlands and Switzerland in boarding schools apart from his mother, who had landed in London, he eventually made it to England.

Ray spent his early English years mostly in boarding schools and actually had rather little time with his mother, who lived in London. He occupied much of his vacation and other times away from school as a houseguest in the home of Wilfrid E. Le Gros Clark, a famous neuroanatomist who by then was the Oxford Professor of Anatomy. Ray's time with Le Gros Clark undoubtedly influenced his ultimate career choice.

#### Higher education

Ray won a scholarship to study medicine at University College London (UCL) from which he received a B.Sc. in 1951. Among his teachers, there were J.Z. Young and Bernard Katz. At some point during his medical training, he realized that he did not want a career as a clinician and rather gravitated to biomedical research. Towards that end, he entered a Ph.D. programme at UCL under the tutelage of J.Z. Young. His main interest centred on the hypothalamus, and his thesis involved a quantitative count of axons in the fornix, a fibre pathway between the hippocampus and mamillary bodies of the hypothalamus. He received his Ph.D. in 1954.

#### Faculty positions

After his Ph.D., Ray stayed on at UCL as a faculty member until 1964, at which time he moved to the USA where he accepted a faculty position in the Department of Anatomy at the University of

Wisconsin. There he founded a neuroscience graduate programme and fostered faculty with interest broadly in neuroscience and visual neuroscience. In 1977, he moved to become Professor in the Department of Pharmacology and Physiology at the University of Chicago. There, he became the founding Chair of the Committee on Neurobiology, thereby starting the first Ph.D.-granting graduate programme devoted to neuroscience at the University of Chicago. In 1984, he returned to England to become Thomas Lee's Professor and Head of the Department of Human Anatomy at the University of Oxford. In 1996, at the age of 65, he moved back to his old Department of Anatomy at the University of Wisconsin with the title of Visiting Professor and Senior Scientist. Then, to be closer to his daughter, who had settled with her husband in Istanbul, Turkey, he moved there in 2006 to become Visiting Professor in the Anatomy Department of Marmara University. Finally, in 2010, he returned to England as an Honorary Emeritus Research Fellow of the MRC Anatomical Neuropharmacology Unit at the University of Oxford. He remained in that position and enjoyed a productive writing period until days before his passing.

### Family life

In 1954, Ray married Margot Pepper, who at the time was completing her medical degree. Four children were born to Ray and Margot during his time at UCL. Peter John Guillery, born on 6 January 1957, is an architectural historian living in London. Nigel Robert (known as Edward) Guillery, born on 18 June 1958, is a paediatric nephrologist living in Portland. Richard Philip (known as Phil) Guillery, born on 25 March 1960, works for sustainable forestry as a Director at the Forest Stewardship Council and lives in Colorado; and Jane Louise Guillery, born on 20 September 1963 and now Jane Kandur, is a teacher and language translator living in Istanbul.

### Scientific contributions

Ray was an extraordinarily productive and insightful scientist. At the beginning of his career, neuroscience was in its infancy as a recognized profession, and the field as such was distributed amongst the more established and recognized divisions at the time, such as anatomy, physiology, psychology, pharmacology. The subdisciplines within neuroscience (e.g., neuroanatomy and neurophysiology) generally ignored one another. However, whereas Ray's early work could be characterized as neuroanatomy, he always had a holistic theoretical framework that informed his approach, so that rather than positioning himself as a descriptive anatomist, Ray always interpreted his findings more broadly and queried their meaning in physiology and behaviour.

Ray trained at UCL with J.Z. Young, and in collaboration with George Gray and Brian Boycott, was the first to describe asymmetrical and symmetrical synapses (often referred to as Gray's type 1 and type 2) using the new technology of electron microscopy. Broadly interested in the cytology and connectivity of neurons, Ray's early work concentrated on hypothalamus and its connections, variously the cytology of the leech nerve cord and of connections in sensory systems of other lower vertebrates (Boycott *et al.*, 1961; Gray & Guillery, 1961, 1963a,b; Boycott & Guillery, 1962). Ray was drawn to and focused on the thalamus and thalamocortical relationships, including development thereof, and used the mammalian visual system as his chief model for investigation. His major contributions can be placed into four main categories: investigation of visual pathways from retina to cortex, investigation of the

development of these pathways, studies of the defect in these pathways associated with albinism, and theoretical considerations of thalamocortical relationships.

### *Studies of retino-geniculo-cortical pathways*

Most of Ray's contributions on this topic were based on the cat model, although other species were also studied. He was an early practitioner of electron microscopy, and he used this to good effect to unpack the circuitry of the lateral geniculate nucleus (Guillery, 1969a,b). His work classified and defined the source of the various synaptic profiles in this thalamic nucleus, and he made the surprising finding that the functionally dominant input from the retina produced surprisingly few synapses onto relay cells. He then extended many of these observations to the monkey (Colonnier & Guillery, 1964; Guillery & Colonnier, 1970). Likewise, in a landmark Golgi study, he defined the classes of relay cells and interneurons in the lateral geniculate nucleus of the cat (Guillery, 1966). Ray also redefined the laminar patterning of the cat's lateral geniculate nucleus (Guillery, 1970; Hickey & Guillery, 1974) and pointed to complexities in lamination of the human lateral geniculate nucleus (Hickey & Guillery, 1979).

### *Studies of retino-geniculo-cortical development*

Ray's developmental studies concentrated on development of the visual pathways in kittens, particularly with respect to the effects of early visual deprivation. Arguably, his most important contribution here was his rigorous demonstration of the role of binocular competition in the development of the visual pathways, that is, competition between pathways related to each eye in the formation of enduring connections in thalamus and cortex. He first showed that cell shrinkage in the deprived layers of the lateral geniculate nucleus in monocularly deprived cats was limited to the binocular segment (Guillery & Stelzner, 1970). This showed that the deleterious effects of deprivation only occurred in regions where the deprived eye would have to compete with the open eye for synaptic connections; in deprived monocular regions, where the deprived eye was by definition at no competitive disadvantage, development proceeded relatively normally.

In an ingenious extension of this, he created an artificial, central monocular segment for the deprived eye by placing a central retinal lesion in the open eye. Again, deprived geniculate cells grew to normal size in the natural and artificial monocular zones of the lateral geniculate nucleus (Guillery, 1972). Finally, in collaboration with one of the authors (SMS), he extended the finding of normal development related to the natural and artificial monocular zones using electrophysiological and behavioural approaches (Sherman *et al.*, 1974, 1975; Sherman & Guillery, 1976).

Starting in the late 1970's, Ray oversaw a swathe of studies on the development of the visual pathways, spanning studies on abnormal sprouting of retinal afferents (Robson *et al.*, 1978), development of layers in the lateral geniculate nucleus (Guillery *et al.*, 1985), and the development of retinogeniculate afferents in the kitten (Mason, 1982a, b). He also analysed retinal representations of axon order in the optic tract (Torrealba *et al.*, 1981, 1982).

### *Studies of visual pathways in albinos*

Albinos have long been known to suffer visual difficulties, but these were usually explained by optical deficiencies: lack of pigment

means that the retinal image is much degraded by light scatter. Ray showed that there was an additional, heretofore unappreciated explanation: retinofugal projections in albinos are aberrant. In the 1970s, Ray demonstrated that albinos of all mammalian species, from Siamese cats to humans to white tigers, show abnormal patterns of nerve crossing at the optic chiasm, with resulting abnormalities in vision, and then discovered how the brain manages this abnormal visual input. He and Jon Kaas first demonstrated abnormalities in connectivity in the Siamese cat (Guillery & Kaas, 1971), which has a thermolabile form of albinism. They demonstrated that a region of geniculate lamina A1, which normally receives retinal input only from the ipsilateral temporal retina, instead is also innervated by the contralateral temporal retina. Ray and colleagues extended this basic observation to a number of other species: tigers (Guillery & Kaas, 1973), mice (Guillery *et al.*, 1973), mink (Sanderson *et al.*, 1974), axolotls (Guillery & Updyke, 1976), ferrets (Cucchiario & Guillery, 1984), monkeys (Guillery *et al.*, 1984) and even humans (Guillery *et al.*, 1975). In an important extension, Ray and Jon Kaas also showed that the defect in albinos affects cortical circuitry as well and thus likely disturbs vision (Kaas & Guillery, 1973). The decussation defect at the optic chiasm of albinos was essentially the first developmental genetic aberration identified in higher mammals and is a paradigm for how genetics can regulate circuit formation and function.

#### *Theoretical considerations of thalamocortical relationships*

Starting in 1996, Ray and one of the authors (SMS) wrote a series of theoretical suggestions for new ways of looking at thalamocortical relationships. This led to 10 review articles and book chapters (Sherman & Guillery, 1996, 1998, 2002, 2004a,b, 2011, 2014; Guillery & Sherman, 2002a,b, 2011) and three monographs (Sherman & Guillery, 2001, 2006, 2013).

Together, they developed three novel ideas. First, they classified glutamatergic pathways in thalamus and cortex as *drivers* or *modulators*, suggesting that only the subset of drivers (e.g., retinal input to geniculate relay cells) carries basic information, whereas modulatory input (e.g., layer 6 corticogeniculate input) serves to modulate the processing of driver input. Second, they identified a novel form of corticocortical communication: layer 5 of a cortical area projects to a thalamic nucleus that relays this input to another cortical area, and these transthalamic circuits connect cortical areas in parallel with direct corticocortical connections. Third, because driver inputs to thalamus often, if not always, arrive via branching axons with the extrathalamic targets often being motor structures in the brainstem, they suggested that these inputs serve as efference copies for motor commands generated by cortex.

The final gift of Ray's broad and important oeuvre is a new book entitled "The Brain as a Tool," the final page proofs of which were received days before he died. This book will be published in fall, 2017, by Oxford University Press. In this monograph, Ray makes an ambitious, and, we believe, successful, attempt to correlate details of thalamocortical circuitry with higher cognitive functions. Ray says it best, so to quote from his preface to the book:

"...this book, and the title is an attempt to see how we depend on using the brain, its nerve cells and neural pathways, to learn about our interactions with the world. The brain on its own can do nothing. We use our brains to interact with the world. It is not until we start moving about, moving our eyes, or our fingers, that we can start using our brains to learn about the world by interacting with it. If we want to understand animals including ourselves and each other in health and disease, we need to understand the neural pathways and their

functions. ... That is the focus of this book: to raise answerable questions about living brains and identifiable intercommunicating nerve cells."

#### Service and Honours

Ray functioned in a number of important service positions. In addition to serving on various editorial boards and grant reviewing committees, Ray was the Editor-in-Chief and founding editor of this journal, *European Journal of Neuroscience*, and a member of editorial boards of *Journal of Neuroscience* (section editor), *Journal of Neurophysiology*, *Journal of Neurocytology*, *Journal of Comparative Neurology*, *Neuroscience*, *Journal of Anatomy*, *Visual Neuroscience*, and *Neuroembryology and Aging*. Ray served as Treasurer of the Society for Neuroscience and President of the Anatomical Society of Great Britain and Ireland. Ray was named as a Fellow of the Royal Society in 1983.

#### Concluding remarks

Ray's gifts to us include his critical eye and intellectual rigour, done with the highest level of scholarship, and often suffused with humour even as he was tough. He let us develop independently, often insisting on excluding his name as author on studies fully done in his laboratory and cultivated by him. He also posed issues and questions, like a latter-day Ramon y Cajal, before he or the field had any knowledge of molecular mechanisms (e.g., Guillery *et al.*, 1995).

Ray's modesty, interest, sharp intellect, humour and selflessness made him unique as a person, a scientist and a mentor. He loved opera, gardening, woodworking, cooking and travelling. He also enjoyed a good dinner with friends during which history, art, philosophy and science were explored and discussed with elegance and humour. During his time in Istanbul, he held long dinners with students and scientists. These meetings helped shape the career paths of many, expanding and adding depth to all subsequent steps taken by them. What made Ray unique was also the way in which he cared for and enabled his mentees to achieve their life goals. He would very kindly allow them to develop their own scientific approaches while gently offering guidance during the process. He had a stunning way of integrating science and life together as a continuum.

As a scientist, he had a remarkable ability to examine deeply the most fundamental aspects of any scientific puzzle, and he worked on explaining one of the most complex puzzles of all time: How does the brain interact with the world around us and generate conscious experiences? We now look forward to reading about this topic in his upcoming book.

He will be greatly missed.

#### References

- Boycott, B.B. & Guillery, R.W. (1962) Olfactory and visual learning in the red-eared terrapin, *pseudemys scripta elegans* (wied.). *J. Exp. Biol.*, **39**, 567–577.
- Boycott, B.B., Gray, E.G. & Guillery, R.W. (1961) Synaptic structure and its alteration with environmental temperature: a study by light and electron microscopy of the central nervous system of lizards. *Proc. R. Soc. Lond B Bio. Sci.*, **154**, 172.
- Colonnier, M. & Guillery, R.W. (1964) Synaptic organization in the lateral geniculate nucleus of the monkey. *Z. Zellforsch.*, **62**, 333–355.
- Cucchiario, J.B. & Guillery, R.W. (1984) The development of the retinogeniculate pathways in normal and albino ferrets. *Proc. R. Soc. Lond. B.*, **223**, 141–164.

- Gray, E.G. & Guillery, R.W. (1961) The basis for silver staining of synapses of the mammalian spinal cord: a light and electron microscope study. *J. Physiol.*, **157**, 581–588.
- Gray, E.G. & Guillery, R.W. (1963a) An electron microscopical study of the ventral nerve cord of the leech. *Z. Zellforsch. Mikrosk. Anat.*, **60**, 826–849.
- Gray, E.G. & Guillery, R.W. (1963b) On nuclear structure in the ventral nerve cord of the leech *Hirudo medicinalis*. *Z. Zellforsch.*, **59**, 738–745.
- Guillery, R.W. (1966) A study of Golgi preparations from the dorsal lateral geniculate nucleus of the adult cat. *J. Comp. Neurol.*, **128**, 21–50.
- Guillery, R.W. (1969a) A quantitative study of synaptic interconnections in the dorsal lateral geniculate nucleus of the cat. *Z. Zellforsch.*, **96**, 39–48.
- Guillery, R.W. (1969b) The organization of synaptic interconnections in the laminae of the dorsal lateral geniculate nucleus of the cat. *Z. Zellforsch.*, **96**, 1–38.
- Guillery, R.W. (1970) The laminar distribution of retinal fibers in the dorsal lateral geniculate nucleus of the cat: A new interpretation. *J. Comp. Neurol.*, **138**, 339–368.
- Guillery, R.W. (1972) Binocular competition in the control of geniculate cell growth. *J. Comp. Neurol.*, **144**, 117–130.
- Guillery, R.W. & Colonnier, M. (1970) Synaptic patterns in the dorsal lateral geniculate nucleus of the monkey. *Z. Zellforsch. Mikrosk. Anat.*, **103**, 90–108.
- Guillery, R.W. & Kaas, J.H. (1971) A study of normal and congenitally abnormal retinogeniculate projections in cats. *J. Comp. Neurol.*, **143**, 73–100.
- Guillery, R.W. & Kaas, J.H. (1973) Genetic abnormality of the visual pathways in a “white” tiger. *Science*, **180**, 1287–1289.
- Guillery, R.W. & Sherman, S.M. (2002a) Thalamic relay functions and their role in corticocortical communication: Generalizations from the visual system. *Neuron*, **33**, 163–175.
- Guillery, R.W. & Sherman, S.M. (2002b) The thalamus as a monitor of motor outputs. *Philos. T. R. Soc. Lon. [B.]*, **357**, 1809–1821.
- Guillery, R.W. & Sherman, S.M. (2011) Branched thalamic afferents: what are the messages that they relay to cortex? *Brain Res. Brain Res. Rev.*, **66**, 205–219.
- Guillery, R.W. & Stelzner, D.J. (1970) The differential effects of unilateral lid closure upon the monocular and binocular segments of the dorsal lateral geniculate nucleus in the cat. *J. Comp. Neurol.*, **139**, 413–422.
- Guillery, R.W. & Updyke, B.V. (1976) Retinofugal pathways in normal and albino axolotls. *Brain Res.*, **109**, 235–244.
- Guillery, R.W., Scott, G.L., Cattanaach, B.M. & Deol, M.S. (1973) Genetic mechanisms determining the central visual pathways of mice. *Science*, **179**, 1014–1016.
- Guillery, R.W., Okoro, A.N. & Witkop, C.J. Jr (1975) Abnormal visual pathways in the brain of a human albino. *Brain Res.*, **96**, 373–377.
- Guillery, R.W., Hickey, T.L., Kaas, J.H., Felleman, D.J., DeBruyn, E.J. & Sparks, D.L. (1984) Abnormal central visual pathways in the brain of an albino green monkey (*Cercopithecus aethiops*). *J. Comp. Neurol.*, **226**, 165–183.
- Guillery, R.W., LaMantia, A.S., Robson, J.A. & Huang, K. (1985) The influence of retinal afferents upon the development of layers in the dorsal lateral geniculate nucleus of mustelids. *J. Neurosci.*, **5**, 1370–1379.
- Guillery, R.W., Mason, C.A. & Taylor, J.S. (1995) Developmental determinants at the mammalian optic chiasm. *J. Neurosci.*, **15**, 4727–4737.
- Hickey, T.L. & Guillery, R.W. (1974) An autoradiographic study of retinogeniculate pathways in the cat and the fox. *J. Comp. Neurol.*, **156**, 239–254.
- Hickey, T.L. & Guillery, R.W. (1979) Variability of laminar patterns in the human lateral geniculate nucleus. *J. Comp. Neurol.*, **183**, 221–246.
- Kaas, J.H. & Guillery, R.W. (1973) The transfer of abnormal visual field representations from the dorsal lateral geniculate nucleus to the visual cortex in siamese cats. *Brain Res.*, **59**, 61–95.
- Mason, C.A. (1982a) Development of terminal arbors of retino-geniculate axons in the kitten—I. *Light microscopical observations. Neuroscience*, **7**, 41–59.
- Mason, C.A. (1982b) Development of terminal arbors of retino-geniculate axons in the kitten—II. *Electron microscopical observations. Neuroscience*, **7**, 561–582.
- Robson, J.A., Mason, C.A. & Guillery, R.W. (1978) Terminal arbors of axons that have formed abnormal connections. *Science*, **201**, 635–637.
- Sanderson, K.J., Guillery, R.W. & Shackelford, R.M. (1974) Congenitally abnormal visual pathways in mink (*Mustela vison*) with reduced retinal pigment. *J. Comp. Neurol.*, **1254**, 225–248.
- Sherman, S.M. & Guillery, R.W. (1976) Behavioral studies of binocular competition in cats. *Vision. Res.*, **16**, 1479–1481.
- Sherman, S.M. & Guillery, R.W. (1996) The functional organization of thalamocortical relays. *J. Neurophysiol.*, **76**, 1367–1395.
- Sherman, S.M. & Guillery, R.W. (1998) On the actions that one nerve cell can have on another: Distinguishing “drivers” from “modulators”. *Proc. Natl. Acad. Sci. USA*, **95**, 7121–7126.
- Sherman, S.M. & Guillery, R.W. (2001) *Exploring the Thalamus*. Academic Press, San Diego.
- Sherman, S.M. & Guillery, R.W. (2002) The role of thalamus in the flow of information to cortex. *Philos. T. R. Soc. Lon. [B.]*, **357**, 1695–1708.
- Sherman, S.M. & Guillery, R.W. (2004a) Thalamus. In Shepherd, G.M. (ed), *Synaptic Organization of the Brain*. Oxford University Press, Oxford, pp. 311–359.
- Sherman, S.M. & Guillery, R.W. (2004b) The visual relays in the thalamus. In Chalupa, L.M. & Werner, J.S. (eds), *The Visual Neurosciences*. MIT Press, Cambridge, MA, pp. 565–591.
- Sherman, S.M. & Guillery, R.W. (2006) *Exploring the Thalamus and its Role in Cortical Function*. MIT Press, Cambridge, MA.
- Sherman, S.M. & Guillery, R.W. (2011) Distinct functions for direct and transthalamic corticocortical connections. *J. Neurophysiol.*, **106**, 1068–1077.
- Sherman, S.M. & Guillery, R.W. (2013) *Thalamocortical Processing: Understanding the Messages that Link the Cortex to the World*. MIT Press, Cambridge, MA.
- Sherman, S.M. & Guillery, R.W. (2014) The lateral geniculate nucleus and pulvinar. In Chalupa, L.M. & Werner, J.S. (Eds), *The New Visual Neurosciences*. MIT Press, Cambridge, MA, pp. 257–283.
- Sherman, S.M., Guillery, R.W., Kaas, J.H. & Sanderson, K.J. (1974) Behavioral, electrophysiological and morphological studies of binocular competition in the development of the geniculo-cortical pathways of cats. *J. Comp. Neurol.*, **158**, 1–18.
- Sherman, S.M., Wilson, J.R. & Guillery, R.W. (1975) Evidence that binocular competition affects the postnatal development of Y-cells in the cat’s lateral geniculate nucleus. *Brain Res.*, **100**, 441–444.
- Torrealba, F., Guillery, R.W., Polley, E.H. & Mason, C.A. (1981) A demonstration of several independent, partially overlapping, retinotopic maps in the optic tract of the cat. *Brain Res.*, **219**, 428–432.
- Torrealba, F., Guillery, R.W., Eysel, U.T., Polley, E.H. & Mason, C.A. (1982) Studies of retinal representations within the cat’s optic tract. *J. Comp. Neurol.*, **211**, 377–396.